

## Report on ENVISAT Radar Altimeter - 2 (RA-2)

### Wind/Wave Product with Height Information (RA2 WWV 2P)

**By:** *Saleh Abdalla*

**Date:** *11 September 2003*

#### **Overview:**

Based on the data received during this month, on average, 9361 (varies between 0 and 19477) observations arrived at ECMWF every 6-hour window of which an average of 3565 observations were rejected initially because of one of the following reasons: being over land, being outside model domain, being a double observation or flagged for rain contamination. On average 78.47% of the remaining part passed the quality control. There was no data during the following periods (in terms of 6-hour time-windows; all times are in UTC):

- Data either missing or significantly reduced at midnight everyday.
- Time windows 18:00 on the 2nd., the 4th., the 6th. and the 10th. of the month.
- Time window 12:00 on the 18th. of the month.
- Time windows 06:00 on the 23rd. and the 26th. of the month.

The time series of the data coverage and some reasons of rejection can be seen in Figure 1. Note that we are talking about the raw data which we downloaded in “bufr” format (since June 2003 bufr data as converted by ESA are used) before they were processed.

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪

## Quality of Received Data:

For the period covered, the RA-2 Ku-band wave height data are generally of good quality. The quality of S-band wave height observations is similar to last month. The S-band wave heights keep their trend to get higher than the Ku-band waves. The quality of wind speed observations is acceptable. Apart from the few outliers, MWR products are generally fine and in good agreement with the model (they still somewhat drier than the model).

## Backscatter:

- ENVISAT RA-2 Ku-Band  $\langle\sigma_0\rangle = 11.17$  dB (with a single peak at  $\sim 11.1$  dB).
- ENVISAT RA-2 S-Band  $\langle\sigma_0\rangle = 10.95$  dB (with a single rather broad peak at  $\sim 10.0$  dB).

## Comparison Summary:

Table 1: Comparison of Surface Wind Speeds:

	RA2 - ECMWF		RA2 - Buoy	
	Bias (m/s)	SI (%)	Bias (m/s)	SI (%)
Global	-0.14	17.8	-0.87	20.7
Northern Hemisphere	-0.62	20.0	-0.88	22.8
Tropics	-0.32	18.2	-0.74	10.5
Southern Hemisphere	0.32	14.7	----	----

# ■ ECMWF Report on ENVISAT RA-2 for August 2003 ■

Table 2: Comparison of Ku-Band Significant Wave Heights:

	RA2 (Ku) - WAM		RA2 (Ku) - Buoy	
	Bias (m)	SI (%)	Bias (m)	SI (%)
Global	0.10	15.7	0.10	16.6
Northern Hemisphere	0.05	20.2	0.11	17.8
Tropics	-0.05	14.7	0.03	8.0
Southern Hemisphere	0.26	12.5	----	----

Table 3: Comparison of S-Band Significant Wave Heights:

	RA2 (S) - WAM		RA2 (S) - Buoy	
	Bias (m)	SI (%)	Bias (m)	SI (%)
Global	0.14	28.8	0.28	35.8
Northern Hemisphere	0.28	46.1	0.31	33.1
Tropics	0.04	34.9	0.10	43.9
Southern Hemisphere	0.13	20.2	----	----

# ■ ECMWF Report on ENVISAT RA-2 for August 2003 ■

Table 4: Comparison of Wet Tropo Correction and Total Column Water Vapour Values:

	MWR WTC - ECMWF WTC		MWR TCWV - ECMWF TCWV	
	Bias (m)	SI (%)	Bias (kg/m <sup>2</sup> )	SI (%)
Global	-0.018	22.0	-0.76	23.1
Northern Hemisphere	-0.013	29.0	-0.15	30.8
Tropics	-0.012	9.8	+0.50	9.5
Southern Hemisphere	-0.026	31.2	-2.19	33.1

## Remarks:

- For the period covered, the RA-2 Ku-band wave height data are generally of good quality. The S-band wave height data are of acceptable quality (with a number of outliers similar to the previous months). The wind speed data are of acceptable quality. The MWR derived quantities are as usual with a number of outliers when compared against the model.
- According to the used land sea mask (which is used for the operational WAM run at ECMWF), more than one third (34% to be exact) of all processed data have been collected over land. This value is too large as the land records are not filtered out.
- Although the rain flag is only responsible for the rejection of 3% of the data this month, it was responsible for rejection of most of the data on the 15th. and the 21st. of the month (lower panel of Figure 1).

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪

- The wind speed data are in good agreement with the wave model and buoy data except for very low wind speeds (below ~ 4 m/s) and for high wind speeds (above 20 m/s).
- **The wind speed algorithm needs some adjustments both in the low wind regime (below ~ 4 m/s) and more importantly for the very high wind regime (above ~ 20 m/s) as can be concluded from the scatter plots in Figures 7-13 and from comparing the histograms in Figures 5 and 6.**
- There is a trend for Ku-band wave heights to be slightly overestimated especially when compared to WAM results by about 6%. This is visually clear in the scatter plots in Figures 22-25 (Ku-band - WAM comparisons) and can be inferred from the symmetric slope values in same scatter plots as well as in Figures 30-32 (Ku-band - buoy comparison). On the other hand, the RA-2 Ku-band wave heights are higher than buoy wave heights by about 4%.
- The S-band wave heights are similar to the previous month when compared to both the WAM and the buoy data with quite a number of outliers when compared to the model (as can be seen in the scatter plots in Figures 26-29) and to the buoy (as can be seen in the scatter plots in Figures 33-34).
- As can be seen in Figure 37, S-band wave heights suffered some degradation on the 2nd. and the 15th. of the month.
- **The S-band wave height values increased, thus the ratio between Ku-band and S-band wave heights decreased, over the period July 2002 to early April 2003 as can be seen in Figure 48. This behaviour was handled on the 9th of April bringing the wave heights from both sensors to a comparable level. However, this lasted for few days only. The S-band wave height keeps their tendency to increase and they are now about 5% higher than the Ku-band wave height as can be clearly seen in Figure 48.**
- There is quite a number of outliers in the scatter plots comparing the MWR derived wet tropospheric correction (WTC) and total column water vapour (TCWV) against the ECMWF model in the Northern and Southern Hemisphere (Figures 39, 41, 43 and 45). The tendency of the ECMWF model to have saturation values for both quantities (around 0.4 m for WTC and around 60 kg/m<sup>2</sup> for TCWV) is still existing. We stress that the model behaviour is theoretically more sound.

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪

- The MWR derived WTC and TCWV tend to be smaller (drier) than ECMWF values for smaller values, then tend to converge towards the ECMWF values at high values.
- It is important to stress that one needs to keep in mind when making the comparison between the results presented here for the ENVISAT RA-2 and the results presented in the ERS-2 altimeter reports that the ERS-2 plots and statistics are done for super-observations composed of 30 individual observation, while the plots and statistics here are for super-observations with 11 individual observations. Therefore, it is natural for the RA-2 plots and statistics to show a bit more variability.
- The ERS-2 radar altimeter and SAR wave data used for assimilation at ECMWF wave model are now limited to the North Atlantic and the western coast of North America following the failure of the ERS-2 tape recorders in June 2003. As a result, the difference statistics between ENVISAT RA-2 and first guess wave model wave heights are slightly higher than those from the previous months. Also note that Figure 19 only represents the histogram for the covered area only.

## Comparison Method:

*A brief description of RA-2 data processing and quality control process together with any special considerations, assumptions and/or data corrections needed while comparing various quantities will go here.*

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪

## List of Figures:

- Figure 1: Time series of data reception for ENVISAT Altimeter data for August 2003.
- Figure 2: Distribution of the ENVISAT Altimeter Ku-band backscatter after QC for August 2003.
- Figure 3: Distribution of the ENVISAT Altimeter S-band backscatter after QC for August 2003.
- Figure 4: Distribution of the ENVISAT Altimeter wind speeds after QC for August 2003.
- Figure 5: Distribution of the ENVISAT Altimeter wind speeds after along track averaging for August 2003.
- Figure 6: Global distribution of ECMWF ocean surface wind speeds for August 2003.
- Figure 7: Comparison between ENVISAT Altimeter and ECMWF surface wind speeds for August 2003 (Global).
- Figure 8: Comparison between ENVISAT Altimeter and ECMWF surface wind speeds for August 2003 (Northern Hemisphere).
- Figure 9: Comparison between ENVISAT Altimeter and ECMWF surface wind speeds for August 2003 (Tropics).
- Figure 10: Comparison between ENVISAT Altimeter and ECMWF surface wind speeds for August 2003 (Southern Hemisphere).
- Figure 11: Comparison between ENVISAT Altimeter and buoy surface wind speeds for August 2003 (Global).
- Figure 12: Comparison between ENVISAT Altimeter and buoy surface wind speeds for August 2003 (Northern Hemisphere).
- Figure 13: Comparison between ENVISAT Altimeter and buoy surface wind speeds for August 2003 (Tropics).
- Figure 14: ENVISAT Altimeter wind speeds: Timeseries of daily bias (RA2 - model) and scatter index for August 2003.
- Figure 15: Distribution of the ENVISAT Altimeter Ku-band wave heights after QC for August 2003.
- Figure 16: Distribution of the ENVISAT Altimeter S-band wave heights after QC for August 2003.
- Figure 17: Distribution of the ENVISAT Altimeter Ku-band wave heights after along track averaging for August 2003.
- Figure 18: Distribution of the ENVISAT Altimeter S-band wave heights after along track averaging for August 2003.
- Figure 19: Distribution of the ERS-2 Altimeter wave heights after along track averaging for August 2003.
- Figure 19b: Distribution of WAM first guess (4V) wave heights collocated with ENVISAT for August 2003.
- Figure 20: Global distribution of WAM first guess wave heights for August 2003.

# ■ ECMWF Report on ENVISAT RA-2 for August 2003 ■

- Figure 21: Global distribution of WAM analysis (ERS-2 RA Assimilation) wave heights for August 2003.
- Figure 22: Comparison between ENVISAT Altimeter Ku-band and WAM significant wave heights for August 2003 (Global).
- Figure 23: Comparison between ENVISAT Altimeter Ku-band and WAM sig. wave heights for August 2003 (Northern Hemisphere).
- Figure 24: Comparison between ENVISAT Altimeter Ku-band and WAM significant wave heights for August 2003 (Tropics).
- Figure 25: Comparison between ENVISAT Altimeter Ku-band and WAM sig. wave heights for August 2003 (Southern Hemisphere).
- Figure 26: Comparison between ENVISAT Altimeter S-band and WAM significant wave heights for August 2003 (Global).
- Figure 27: Comparison between ENVISAT Altimeter S-band and WAM sig. wave heights for August 2003 (Northern Hemisphere).
- Figure 28: Comparison between ENVISAT Altimeter S-band and WAM significant wave heights for August 2003 (Tropics).
- Figure 29: Comparison between ENVISAT Altimeter S-band and WAM sig. wave heights for August 2003 (Southern Hemisphere).
- Figure 30: Comparison between ENVISAT Altimeter Ku-band and buoy significant wave heights for August 2003 (Global).
- Figure 31: Comparison between ENVISAT Altimeter Ku-band and buoy sig. wave heights for August 2003 (Northern Hemisphere).
- Figure 32: Comparison between ENVISAT Altimeter Ku-band and buoy significant wave heights for August 2003 (Tropics).
- Figure 33: Comparison between ENVISAT Altimeter S-band and buoy significant wave heights for August 2003 (Global).
- Figure 34: Comparison between ENVISAT Altimeter S-band and buoy sig. wave heights for August 2003 (Northern Hemisphere).
- Figure 35: Comparison between ENVISAT Altimeter S-band and buoy significant wave heights for August 2003 (Tropics).
- Figure 36: ENVISAT Altimeter Ku-band wave heights: Timeseries of daily bias (RA2 - model) and scatter index for August 2003.
- Figure 37: ENVISAT Altimeter S-band wave heights: Timeseries of daily bias (RA2 - model) and scatter index for August 2003.
- Figure 38: Comparison between ENVISAT MWR and ECMWF wet tropospheric correction for August 2003 (Global).
- Figure 39: Comparison between ENVISAT MWR and ECMWF wet tropospheric correction for August 2003 (Northern Hemisphere).
- Figure 40: Comparison between ENVISAT MWR and ECMWF wet tropospheric correction for August 2003 (Tropics).
- Figure 41: Comparison between ENVISAT MWR and ECMWF wet tropospheric correction for August 2003 (Southern Hemisphere).
- Figure 42: Comparison between ENVISAT MWR and ECMWF total column water vapour for August 2003 (Global).
- Figure 43: Comparison between ENVISAT MWR and ECMWF total column water vapour for August 2003 (Northern Hemisphere).
- Figure 44: Comparison between ENVISAT MWR and ECMWF total column water vapour for August 2003 (Tropics).

# ■ ECMWF Report on ENVISAT RA-2 for August 2003 ■

Figure 45: Comparison between ENVISAT MWR and ECMWF total column water vapour for August 2003 (Southern Hemisphere).

Figure 46: ENVISAT MWR wet tropospheric correction: Timeseries of daily bias (MWR-model) and scatter index for August 2003.

Figure 47: ENVISAT MWR total column water vapour: Timeseries of daily bias (MWR-model) and scatter index for August 2003.

Figure 48. Timeseries of daily global ratio between mean Ku-Band to mean S-Band significant wave heights since the 18th. of July 2002.

# ■ ECMWF Report on ENVISAT RA-2 for August 2003 ■




Figure 1: Time series of data reception for ENVISAT Altimeter data for August 2003

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 2: Distribution of the ENVISAT Altimeter Ku Backscatter after QC for August 2003

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 3: Distribution of the ENVISAT Altimeter S Backscatter after QC for August 2003

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 4: Distribution of the ENVISAT Altimeter Wind Speed after QC for August 2003

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 5: Distribution of ENVISAT Altimeter Wind Speeds after Along-Track Averaging for August 2003

Saleh Abdalla

European Centre for Medium Range Weather Forecasts  
Shinfield Park, Reading, Berkshire RG2 9AX, England  
Telephone: U.K. (0118) 949 9703, International (+44 118) 949 9703  
Telex 984 7908 ECMWF G, Telefax (0118) 986 9450, e-mail abdalla@ecmwf.int

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 6: Global distribution of ECMWF Analysis ocean surface wind speeds for August 2003

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 7. Comparison between ENVISAT Altimeter and ECMWF wind speeds for August 2003 (Global)

## STATISTICS

ENTRIES	53589
MEAN ECMWF	7.4100
MEAN ENVISAT	7.2678
BIAS (ENVISAT - ECMWF)	- .1423
STANDARD DEVIATION	1.3152
SCATTER INDEX	.1775
CORRELATION	.9429
SYMMETRIC SLOPE	1.0097 ( .0017)
REGR. COEFFICIENT	1.0709 ( .0016)
REGR. CONSTANT	- .6674 ( .0133)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 8. Comparison between ENVISAT Altimeter and ECMWF wind speeds for August 2003 (N.Hem.)

# ■ ECMWF Report on ENVISAT RA-2 for August 2003 ■




Figure 9. Comparison between ENVISAT Altimeter and ECMWF wind speeds for August 2003 (Tropics)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 10. Comparison between ENVISAT Altimeter and ECMWF wind speeds for August 2003 (S.Hem.)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 11. Comparison between ENVISAT Altimeter and buoy wind speeds for August 2003 (Global)

## STATISTICS

ENTRIES	581
MEAN BUOY	6.7260
MEAN ENVISAT	5.8582
BIAS (ENVISAT - BUOY)	-.8678
STANDARD DEVIATION	1.3946
SCATTER INDEX	.2073
CORRELATION	.8761
SYMMETRIC SLOPE	.8990 ( .0214)
REGR. COEFFICIENT	.9249 ( .0211)
REGR. CONSTANT	-.3630 ( .1534)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪





Figure 12. Comparison between ENVISAT Altimeter and buoy wind speeds for August 2003 (N.Hem.)



## STATISTICS

ENTRIES	490
MEAN BUOY	6.4311
MEAN ENVISAT	5.5511
BIAS (ENVISAT - BUOY)	- .8800
STANDARD DEVIATION	1.4684
SCATTER INDEX	.2283
CORRELATION	.8675
SYMMETRIC SLOPE	.8970 ( .0243)
REGR. COEFFICIENT	.9229 ( .0240)
REGR. CONSTANT	- .3839 ( .1675)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 13. Comparison between ENVISAT Altimeter and buoy wind speeds for August 2003 (Tropics)

## STATISTICS

ENTRIES	85
MEAN BUOY	8.3960
MEAN ENVISAT	7.6528
BIAS (ENVISAT - BUOY)	- .7432
STANDARD DEVIATION	.8816
SCATTER INDEX	.1050
CORRELATION	.8731
SYMMETRIC SLOPE	.9128 ( .0514)
REGR. COEFFICIENT	.8219 ( .0504)
REGR. CONSTANT	.7520 ( .4324)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 14: ENVISAT Altimeter wind speeds: Timeseries of bias (ENVISAT - ECMWF) and scatter index (SI)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪





Figure 15: Distribution of the ENVISAT Altimeter Ku Wave Height after QC for August 2003

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪



# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 17: Distribution of ENVISAT Altimeter Ku-Band Wave Heights after Along-Track Averaging for August 2003

Saleh Abdalla

European Centre for Medium Range Weather Forecasts  
Shinfield Park, Reading, Berkshire RG2 9AX, England  
Telephone: U.K. (0118) 949 9703, International (+44 118) 949 9703  
Telex 984 7908 ECMWF G, Telefax (0118) 986 9450, e-mail abdalla@ecmwf.int

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 18: Distribution of ENVISAT Altimeter S-Band Wave Heights after Along-Track Averaging for August 2003

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 19: Distribution of ERS-2 Altimeter Wave Heights after Along-Track Averaging for August 2003

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 19b: Distribution of WAM 4V Wave Height (Collocated with ENVISAT) for August 2003

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 20: Global distribution of ECMWF First-Guess wave heights for August 2003

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 21: Global distribution of ECMWF Analysis (using ERS-2 data) wave heights for August 2003

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 22. Comparison between ENVISAT Altimeter Ku-Band and WAM (first guess) significant wave heights for August 2003 (Global)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 23. Comparison between ENVISAT Altimeter Ku-Band and WAM (first guess) significant wave heights for August 2003 (N.Hem.)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 24. Comparison between ENVISAT Altimeter Ku-Band and WAM (first guess) significant wave heights for August 2003 (Tropics)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 25. Comparison between ENVISAT Altimeter Ku-Band and WAM (first guess) significant wave heights for August 2003 (S.Hem.)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 26. Comparison between ENVISAT Altimeter S-Band and WAM (first guess) significant wave heights for August 2003 (Global)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 27. Comparison between ENVISAT Altimeter S-Band and WAM (first guess) significant wave heights for August 2003 (N.Hem.)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 28. Comparison between ENVISAT Altimeter S-Band and WAM (first guess) significant wave heights for August 2003 (Tropics)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪





Figure 29. Comparison between ENVISAT Altimeter S-Band and WAM (first guess) significant wave heights for August 2003 (S.Hem.)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪



## STATISTICS

ENTRIES	1001
MEAN BUOY	1.5273
MEAN ENVISAT	1.6285
BIAS (ENVISAT - BUOY)	.1012
STANDARD DEVIATION	.2535
SCATTER INDEX	.1660
CORRELATION	.9374
SYMMETRIC SLOPE	1.0385 (.0118)
REGR. COEFFICIENT	.8477 (.0100)
REGR. CONSTANT	.3339 (.0168)

Figure 30. Comparison between ENVISAT Altimeter Ku-Band and buoy significant wave heights for August 2003 (Global)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 31. Comparison between ENVISAT Altimeter Ku-Band and buoy significant wave heights for August 2003 (N.Hem.)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 32. Comparison between ENVISAT Altimeter Ku-Band and buoy significant wave heights for August 2003 (Tropics)

## STATISTICS

ENTRIES	105
MEAN BUOY	2.0417
MEAN ENVISAT	2.0766
BIAS (ENVISAT - BUOY)	.0349
STANDARD DEVIATION	.1624
SCATTER INDEX	.0795
CORRELATION	.9140
SYMMETRIC SLOPE	1.0164 ( .0412)
REGR. COEFFICIENT	.9105 ( .0398)
REGR. CONSTANT	.2176 ( .0828)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 33. Comparison between ENVISAT Altimeter S-Band and buoy significant wave heights for August 2003 (Global)

## STATISTICS

ENTRIES	991
MEAN BUOY	1.5317
MEAN ENVISAT	1.8158
BIAS (ENVISAT - BUOY)	.2841
STANDARD DEVIATION	.5476
SCATTER INDEX	.3575
CORRELATION	.6691
SYMMETRIC SLOPE	1.1247 ( .0270)
REGR. COEFFICIENT	.5349 ( .0189)
REGR. CONSTANT	.9966 ( .0321)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 34. Comparison between ENVISAT Altimeter S-Band and buoy significant wave heights for August 2003 (N.Hem.)

## STATISTICS

ENTRIES	879
MEAN BUOY	1.4726
MEAN ENVISAT	1.7791
BIAS (ENVISAT - BUOY)	.3065
STANDARD DEVIATION	.4869
SCATTER INDEX	.3307
CORRELATION	.7502
SYMMETRIC SLOPE	1.1247 ( .0263)
REGR. COEFFICIENT	.5196 ( .0155)
REGR. CONSTANT	1.0139 ( .0255)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 35. Comparison between ENVISAT Altimeter S-Band and buoy significant wave heights for August 2003 (Tropics)

## STATISTICS

ENTRIES	105
MEAN BUOY	2.0417
MEAN ENVISAT	2.1430
BIAS (ENVISAT - BUOY)	.1013
STANDARD DEVIATION	.8967
SCATTER INDEX	.4392
CORRELATION	.3230
SYMMETRIC SLOPE	1.1254 ( .2301)
REGR. COEFFICIENT	.7762 ( .2241)
REGR. CONSTANT	.5582 ( .4659)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 36: ENVISAT Altimeter Ku-band wave heights: Timeseries of bias (ENVISAT - WAM\_FG) and scatter index (SI)

# ■ ECMWF Report on ENVISAT RA-2 for August 2003 ■




Figure 37: ENVISAT Altimeter S-band wave heights: Timeseries of bias (ENVISAT - WAM\_FG) and scatter index (SI)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 38. Comparison between ENVISAT MWR and ECMWF (analysis) wet tropo correction for August 2003 (Global)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 39. Comparison between ENVISAT MWR and ECMWF (analysis) wet tropo correction for August 2003 (N.Hem.)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪





Figure 40. Comparison between ENVISAT MWR and ECMWF (analysis) wet tropo correction for August 2003 (Tropics)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪



## STATISTICS

ENTRIES	20986
MEAN ECMWF	.0923
MEAN ENVISAT	.0668
BIAS (ENVISAT - ECMWF)	-.0255
STANDARD DEVIATION	.0288
SCATTER INDEX	.3115
CORRELATION	.8587
SYMMETRIC SLOPE	.8362 ( .0045)
REGR. COEFFICIENT	.9908 ( .0041)
REGR. CONSTANT	-.0246 ( .0004)

Figure 41. Comparison between ENVISAT MWR and ECMWF (analysis) wet tropo correction for August 2003 (S.Hem.)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 42. Comparison between ENVISAT MWR and ECMWF (analysis) total column water vapour for August 2003 (Global)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 43. Comparison between ENVISAT MWR and ECMWF (analysis) total column water vapour for August 2003 (N.Hem.)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 44. Comparison between ENVISAT MWR and ECMWF (analysis) total column water vapour for August 2003 (Tropics)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 45. Comparison between ENVISAT MWR and ECMWF (analysis) total column water vapour for August 2003 (S.Hem.)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 46: ENVISAT Altimeter wet tropo correction: Timeseries of bias (ENVISAT - ECMWF) and scatter index (SI)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 47: ENVISAT Altimeter total column water vapour: Timeseries of bias (ENVISAT - ECMWF) and scatter index (SI)

# ▪ ECMWF Report on ENVISAT RA-2 for August 2003 ▪




Figure 48. Timeseries of daily global ratio between mean Ku-Band to mean S-Band significant wave heights since the 18th. of July 2002.