

▪ ECMWF Report on ERS-2 RA for April 2001 ▪

Title: Report on ERS-2 Radar Altimeter wave height and wind speed data.

By: Saleh Abdalla

Date: 8 May 2001

Overview:

Based on the data received during the full month, on average, 12160 observations arrived at ECMWF every 6 hours of which 81.6% passed the quality control for this month. The data coverage can be seen in Figure 1. Note that we are talking about the raw data which have arrived at ECMWF before they were processed.

The quality of the received data, especially that of the wind speed in the southern hemisphere, seems to be improved compared to previous months. This can be clearly recognized when comparing Figure 9 and the scatter plots for this month with those of previous months.

Backscatter:

ERS-2 $\langle\sigma_0\rangle = 10.86 \text{ dB}$ (with two main peaks at 10.3 dB and 11.1 dB)

▪ ECMWF Report on ERS-2 RA for April 2001 ▪

Wind Speed Comparison with ECMWF wind speeds (bias):

ERS-2 global: 0.425 m/s

ERS-2 northern hemisphere: 0.250 m/s

ERS-2 tropics: 0.213 m/s

ERS-2 southern hemisphere: 0.649 m/s

Wind Speed Comparison with buoy wind speeds (bias):

ERS-2 global: -0.197 m/s

ERS-2 northern hemisphere: -0.185 m/s

ERS-2 tropics: -0.283m/s

Wave Height Comparison with ECMWF wave heights (bias):

ERS-2 global: -0.044 m (lowest waves measured: 0.6m)

ERS-2 northern hemisphere: 0.027 m

ERS-2 tropics: -0.074 m

ERS-2 southern hemisphere: -0.059 m

Wave Height Comparison with buoy wave heights (bias):

ERS-2 global: -0.11 m

ERS-2 northern hemisphere: -0.11 m

ERS-2 tropics: -0.20 m

▪ ECMWF Report on ERS-2 RA for April 2001 ▪

Remarks:

- The Altimeter data quality seems to be as good as that before January 2001 when we start having the “Sun Blinding Effect” and, later on, the switching to the Extra Backup Mode (EBM).

Comparison Method:

The Altimeter wave height and wind speed data, as received by ECMWF from ESA through GTS, are the so-called fast delivery products. At ECMWF these data are subject to a quality control method, the details of which are described by Janssen et al. (1989) and Bauer et al. (1992). Consequently, superobservations are formed by averaging 30 consecutive data in order to match the spatial scales of the operational WAM model. Therefore, the collocation statistics are based on the comparison between these superobservations and operational wavemodel products.

In addition, since also wave observations from buoys are received through the GTS, the Altimeter products are also compared against buoy observations. Again, in order to have matching scales, the buoy observations are averaged over a six hour time window. Apart from this, also a height correction is applied to the wind speed observations, since not all buoys observe the winds at the standard height of 10 m. A default observation height of 5 m is assumed, and when available the actual observation height is used. In order to interpolate from the observation height to the standard height a logarithmic wind profile with a roughness length as given by the Charnock relation is assumed, where the Charnock parameter is given the constant value of 0.018.

▪ ECMWF Report on ERS-2 RA for April 2001 ▪

Figure captions:

- Figure 1: Time series of data reception for ERS-2 Altimeter data for April 2001.
- Figure 2: Distribution of the ERS-2 Altimeter Backscatter after QC for April 2001.
- Figure 3: Distribution of the ERS-2 Altimeter wind speeds after QC for April 2001.
- Figure 4: Distribution of the ERS-2 Altimeter wind speeds after along track averaging for April 2001.
- Figure 5: Global distribution of ECMWF ocean surface wind speeds for April 2001.
- Figure 6: Comparison of ECMWF wind speed results with ERS-2 Altimeter wind speed data for April 2001 (global).
- Figure 7: Comparison of ECMWF wind speed results with ERS-2 Altimeter wind speed data for April 2001 (northern hemisphere)
- Figure 8: Comparison of ECMWF wind speed results with ERS-2 Altimeter wind speed data for April 2001 (tropics)
- Figure 9: Comparison of ECMWF wind speed results with ERS-2 Altimeter wind speed data for April 2001 (southern hemisphere)
- Figure 10: Comparison of buoy wind speed observations with ERS-2 Altimeter wind speed data for April 2001 (global).
- Figure 11: Comparison of buoy wind speed observations with ERS-2 Altimeter wind speed data for April 2001 (northern hemisphere).
- Figure 12: Comparison of buoy wind speed observations with ERS-2 Altimeter wind speed data for April 2001 (tropics).
- Figure 13: ERS-2 Altimeter wind speeds: Timeseries of bias (ERS-2 - model) and scatter index (SI).
- Figure 14: Distribution of the ERS-2 Altimeter wave heights after QC for April 2001.
- Figure 15: Distribution of the ERS-2 Altimeter wave heights after along track averaging for April 2001.
- Figure 16: Global distribution of ECMWF wave heights for April 2001.
- Figure 17: Comparison of ECMWF wave height results with ERS-2 Altimeter wave height data for April 2001 (global).
- Figure 18: Comparison of ECMWF wave height results with ERS-2 Altimeter wave height data for April 2001 (northern hemisphere)
- Figure 19: Comparison of ECMWF wave height results with ERS-2 Altimeter wave height data for April 2001 (tropics)
- Figure 20: Comparison of ECMWF wave height results with ERS-2 Altimeter wave height data for April 2001 (southern hemisphere)
- Figure 21: Comparison of buoy wave height observations with ERS-2 Altimeter wave height data for April 2001 (global).
- Figure 22: Comparison of buoy wave height observations with ERS-2 Altimeter wave height data for April 2001 (northern hemisphere).

▪ ECMWF Report on ERS-2 RA for April 2001 ▪

Figure 23: Comparison of buoy wave height observations with ERS-2 Altimeter wave height data for April 2001 (tropics).

Figure 24: ERS-2 Altimeter wave heights: Timeseries of bias (ERS-2 - model) and scatter index (SI) for April 2001.

Figure 25: ERS-2 Altimeter wave heights: Timeseries of bias (ERS-2 - model) and scatter index (SI) from December 1996 to April 2001

Figure 26: ERS-2 Altimeter wind speeds: Timeseries of bias (ERS-2 - model) and scatter index (SI) from December 1996 to April 2001

■ ECMWF Report on ERS-2 RA for April 2001 ■




Figure 1: Time series of data reception for ERS-2 Altimeter data for April 2001

Saleh Abdalla

European Centre for Medium Range Weather Forecasts
Shinfield Park, Reading, Berkshire RG2 9AX, England
Telephone: U.K. (0118) 949 9703, International (+44 118) 949 9703
Telex 984 7908 ECMWF G, Telefax (0118) 986 9450, e-mail: abdalla@ecmwf.int

■ ECMWF Report on ERS-2 RA for April 2001 ■




Figure 2: Distribution of the ERS-2 Altimeter backscatter after QC for April 2001

■ ECMWF Report on ERS-2 RA for April 2001 ■




Figure 3: Distribution of the ERS-2 Altimeter wind speeds after QC for April 2001

■ ECMWF Report on ERS-2 RA for April 2001 ■




Figure 4: Distribution of ERS-2 Altimeter wind speeds after along track averaging for April 2001

■ ECMWF Report on ERS-2 RA for April 2001 ■




Figure 5: Global distribution of ECMWF ocean surface wind speeds for April 2001

▪ ECMWF Report on ERS-2 RA for April 2001 ▪




Figure 6. Comparison of ECMWF wind speed results with ERS2 Altimeter wind speed data for April 2001 (global)

▪ ECMWF Report on ERS-2 RA for April 2001 ▪




Figure 7. Comparison of ECMWF wind speed results with ERS2 Altimeter wind speed data for April 2001 (n.hem.)

▪ ECMWF Report on ERS-2 RA for April 2001 ▪




Figure 8. Comparison of ECMWF wind speed results with ERS2 Altimeter wind speed data for April 2001 (tropics)

■ ECMWF Report on ERS-2 RA for April 2001 ■




Figure 9. Comparison of ECMWF wind speed results with ERS2 Altimeter wind speed data for April 2001 (s.hem.)

▪ ECMWF Report on ERS-2 RA for April 2001 ▪




Figure 10. Comparison of buoy wind speed observations with ERS2 Altimeter wind speed data for April 2001 (global)

▪ ECMWF Report on ERS-2 RA for April 2001 ▪




Figure 11. Comparison of buoy wind speed observations with ERS2 Altimeter wind speed data for April 2001 (n.hem.)

▪ ECMWF Report on ERS-2 RA for April 2001 ▪




Figure 12. Comparison of buoy wind speed observations with ERS2 Altimeter wind speed data for April 2001 (hawaii)

▪ ECMWF Report on ERS-2 RA for April 2001 ▪




Figure 13: ERS-2 Altimeter wind speeds: Timeseries of bias (ERS-2 - model) and scatter index (SI)

■ ECMWF Report on ERS-2 RA for April 2001 ■




Figure 14: Distribution of the ERS-2 Altimeter wave heights after QC for April 2001

■ ECMWF Report on ERS-2 RA for April 2001 ■




Figure 15: Distribution of ERS-2 Altimeter wave heights after along track averaging for April 2001

■ ECMWF Report on ERS-2 RA for April 2001 ■




Figure 16: Global distribution of ECMWF wave heights for April 2001

■ ECMWF Report on ERS-2 RA for April 2001 ■




Figure 17. Comparison of ECMWF wave height results with ERS2 Altimeter wave height data for April 2001 (global)

▪ ECMWF Report on ERS-2 RA for April 2001 ▪




Figure 18. Comparison of ECMWF wave height results with ERS2 Altimeter wave height data for April 2001 (n.hem.)

▪ ECMWF Report on ERS-2 RA for April 2001 ▪




Figure 19. Comparison of ECMWF wave height results with ERS2 Altimeter wave height data for April 2001 (tropics)

▪ ECMWF Report on ERS-2 RA for April 2001 ▪




Figure 20. Comparison of ECMWF wave height results with ERS2 Altimeter wave height data for April 2001 (s.hem.)

▪ ECMWF Report on ERS-2 RA for April 2001 ▪




Figure 21. Comparison of buoy wave height observations with ERS2 Altimeter wave height data for April 2001 (global)

▪ ECMWF Report on ERS-2 RA for April 2001 ▪




Figure 22. Comparison of buoy wave height observations with ERS2 Altimeter wave height data for April 2001 (n.hem.)

▪ ECMWF Report on ERS-2 RA for April 2001 ▪




Figure 23. Comparison of buoy wave height observations with ERS2 Altimeter wave height data for April 2001 (hawaii)

▪ ECMWF Report on ERS-2 RA for April 2001 ▪




Figure 24: ERS-2 Altimeter wave heights: Timeseries of bias (ERS-2 - model) and scatter index (SI)

▪ ECMWF Report on ERS-2 RA for April 2001 ▪




Figure 25: ERS-2 Altimeter wave heights: Timeseries of bias (ERS-2 - model) and scatter index (SI)

▪ ECMWF Report on ERS-2 RA for April 2001 ▪




Figure 26: ERS-2 Altimeter wind speeds: Timeseries of bias (ERS-2 - model) and scatter index (SI)