

▪ ECMWF Report on ERS-2 RA for January 2010 ▪

Title: Report on ERS-2 Radar Altimeter wave height and wind speed data.

By: Saleh Abdalla

Date: 9 February 2010

Overview:

Based on the data received at ECMWF during the whole month, on average, 3147 observations arrived at ECMWF every 6 hours of which 70.37% passed the quality control. As can be seen in Figure 1, there was no data during the 6-hour time windows centred at (all times are UTC and all windows are inclusive):

- from 06:00 on the 2nd to 18:00 on the 3rd,
- from 18:00 on the 4th to 12:00 on the 5th,
- from 18:00 on the 24th to 12:00 on the 25th, and
- from 12:00 on the 30th to 18:00 on the 31st (and continues into next month) of the month.

The current coverage of the data covers: the North Atlantic, the western coast of North America, the eastern and southeastern coasts of Asia, the Southern Ocean and around South Africa (recently no data arrived from this station) as can be seen in Figure 27. The quality of the received data is as good as usual.

Backscatter:

ERS-2 $\langle s_0 \rangle = 10.89$ dB (with a main peak at 11.1 dB).

■ ECMWF Report on ERS-2 RA for January 2010 ■

Wind Speed Comparison with ECMWF wind speeds (bias):

ERS-2 global: +0.151 m/s

ERS-2 northern hemisphere: +0.386 m/s

ERS-2 tropics: -0.283 m/s

ERS-2 southern hemisphere: +0.141 m/s

Wind Speed Comparison with buoy wind speeds (bias):

ERS-2 global: +0.109 m/s

ERS-2 northern hemisphere: +0.390 m/s

ERS-2 tropics (Hawaii): *missing*

Wave Height Comparison with ECMWF wave heights (bias):

ERS-2 global: -0.046 m (lowest waves measured: 0.6m)

ERS-2 northern hemisphere: -0.086 m

ERS-2 tropics: +0.040 m

ERS-2 southern hemisphere: -0.123 m

Wave Height Comparison with buoy wave heights (bias):

ERS-2 global: -0.079 m

ERS-2 northern hemisphere: -0.090 m

ERS-2 tropics (Hawaii): *missing*

▪ ECMWF Report on ERS-2 RA for January 2010 ▪

Remarks:

- The quality of Altimeter data, both wind speed and significant wave height, is nominal.
- Note that the large fluctuations in the Southern Hemispheric (and Tropical) bias in Figures 13 and 24 are basically due to the limited number of daily measurements.
- According to ESA (EOHelp message of 4 July 2003), the recording capabilities of ERS-2 are declared permanently unavailable following the failure of the ERS-2 tape recorder on 22 June 2003. The ERS-2 tape recorders were used to record the ERS-2 Low Rate mission globally for a period of 8 years of continuous acquisition. The global coverage is, therefore, discontinued. The ERS-2 Low Rate mission will be continued within the visibility of ESA ground stations over Europe, North Atlantic, the Arctic and western North America. Coverage extended to include eastern coast of China and around Korea and Japan since 25 June 2005, the Southern Ocean since 4 July 2005, the north eastern parts of the Indian Ocean since 19 October 2006 and around South Africa since 23 May 2008 (no data was received recently from this station). Current coverage can be seen in Figure 27.
- Assimilation of ERS-2 RA wave heights into ECMWF wave model was stopped on 21st. of October 2003 and replaced by ENVISAT RA-2 Ku-band wave heights. Both instruments can not be assimilated at the same time as both satellites follow the same track with relatively short time separation (both can be assimilated safely if they were not so close).
- ENVISAT RA-2 Ku-band significant wave height and ASAR Wave Mode Level 1b data as well as Jason-1 and Jason-2 altimeter significant wave height data are assimilated in the ECMWF wave model.
- The horizontal resolution of ECMWF models was upgraded on the 26th of January 2010. The IFS runs on T1279 which is equivalent to ~16 km (N640). The global wave model now runs on irregular grid of 0.25° (equivalent to ~28 km) using a spectral grid of 36 frequencies and 36 directions. The current operational IFS cycle is CY36R1 (since the 26th of January 2010).

▪ ECMWF Report on ERS-2 RA for January 2010 ▪

Comparison Method:

The Altimeter wave height and wind speed data, as received by ECMWF from ESA through GTS, are the so-called fast delivery products. At ECMWF these data are subject to a quality control method, the details of which are described by Janssen et al. (1989) and Bauer et al. (1992). Consequently, superobservations are formed by averaging 30 consecutive data in order to match the spatial scales of the operational WAM model. Therefore, the collocation statistics are based on the comparison between these superobservations and operational wavemodel products.

In addition, since also wave observations from buoys are received through the GTS, the Altimeter products are also compared against buoy observations. Again, in order to have matching scales, the buoy observations are averaged over a six hour time window. Apart from this, also a height correction is applied to the wind speed observations, since not all buoys observe the winds at the standard height of 10 m. A default observation height of 5 m is assumed, and when available the actual observation height is used. In order to interpolate from the observation height to the standard height a logarithmic wind profile with a roughness length as given by the Charnock relation is assumed, where the Charnock parameter is given the constant value of 0.018.

Figure captions:

Figure 1: Time series of data reception for ERS-2 Altimeter data for January 2010.

Figure 2: Distribution of the ERS-2 Altimeter Backscatter after QC for January 2010.

Figure 3: Distribution of the ERS-2 Altimeter wind speeds after QC for January 2010.

Figure 4: Distribution of the ERS-2 Altimeter wind speeds after along track averaging for January 2010.

Figure 5: Global distribution of ECMWF ocean surface wind speeds for January 2010.

Figure 6: Comparison of ECMWF wind speed results with ERS-2 Altimeter wind speed data for January 2010 (global).

■ ECMWF Report on ERS-2 RA for January 2010 ■

- Figure 7: Comparison of ECMWF wind speed results with ERS-2 Altimeter wind speed data for January 2010 (northern hemisphere)
- Figure 8: Comparison of ECMWF wind speed results with ERS-2 Altimeter wind speed data for January 2010 (tropics)
- Figure 9: Comparison of ECMWF wind speed results with ERS-2 Altimeter wind speed data for January 2010 (southern hemisphere)
- Figure 10: Comparison of buoy wind speed observations with ERS-2 Altimeter wind speed data for January 2010 (global).
- Figure 11: Comparison of buoy wind speed observations with ERS-2 Altimeter wind speed data for January 2010 (northern hemisphere).
- Figure 12: Comparison of buoy wind speed observations with ERS-2 Altimeter wind speed data for January 2010 (hawaii).
- Figure 13: ERS-2 Altimeter wind speeds: Timeseries of bias (ERS-2 - model) and scatter index (SI).
- Figure 14: Distribution of the ERS-2 Altimeter wave heights after QC for January 2010.
- Figure 15: Distribution of the ERS-2 Altimeter wave heights after along track averaging for January 2010.
- Figure 16: Global distribution of ECMWF wave heights for January 2010.
- Figure 17: Comparison of ECMWF wave height results with ERS-2 Altimeter wave height data for January 2010 (global).
- Figure 18: Comparison of ECMWF wave height results with ERS-2 Altimeter wave height data for January 2010 (northern hemisphere)
- Figure 19: Comparison of ECMWF wave height results with ERS-2 Altimeter wave height data for January 2010 (tropics)
- Figure 20: Comparison of ECMWF wave height results with ERS-2 Altimeter wave height data for January 2010 (southern hemisphere)
- Figure 21: Comparison of buoy wave height observations with ERS-2 Altimeter wave height data for January 2010 (global).
- Figure 22: Comparison of buoy wave height observations with ERS-2 Altimeter wave height data for January 2010 (northern hemisphere).
- Figure 23: Comparison of buoy wave height observations with ERS-2 Altimeter wave height data for January 2010 (hawaii).
- Figure 24: ERS-2 Altimeter wave heights: Timeseries of bias (ERS-2 - model) and scatter index (SI) for January 2010.
- Figure 25: ERS-2 Altimeter wave heights: Timeseries of bias (ERS-2 - model) and scatter index (SI) from December 1996 to January 2010.
- Figure 26: ERS-2 Altimeter wind speeds: Timeseries of bias (ERS-2 - model) and scatter index (SI) from December 1996 to January 2010.
- Figure 27: Significant wave height: Monthly mean difference of ERS-2 altimeter data minus wave model results for January 2010.

■ ECMWF Report on ERS-2 RA for January 2010 ■




Figure 1: Time series of data reception for ERS-2 Altimeter data for January 2010

▪ ECMWF Report on ERS-2 RA for January 2010 ▪




Figure 2: Distribution of the ERS-2 Altimeter backscatter after QC for January 2010

■ ECMWF Report on ERS-2 RA for January 2010 ■




Figure 3: Distribution of the ERS-2 Altimeter wind speeds after QC for January 2010

■ ECMWF Report on ERS-2 RA for January 2010 ■




Figure 4: Distribution of ERS-2 Altimeter wind speeds after along track averaging for January 2010

■ ECMWF Report on ERS-2 RA for January 2010 ■




Figure 5: Global distribution of ECMWF ocean surface wind speeds for January 2010

ECMWF Report on ERS-2 RA for January 2010




Figure 6. Comparison of ECMWF wind speed results with ERS2 Altimeter wind speed data for January 2010 (global)

ECMWF Report on ERS-2 RA for January 2010




Figure 7. Comparison of ECMWF wind speed results with ERS2 Altimeter wind speed data for January 2010 (n.hem.)

ECMWF Report on ERS-2 RA for January 2010




Figure 8. Comparison of ECMWF wind speed results with ERS2 Altimeter wind speed data for January 2010 (tropics)

■ ECMWF Report on ERS-2 RA for January 2010 ■




Figure 9. Comparison of ECMWF wind speed results with ERS2 Altimeter wind speed data for January 2010 (s.hem.)

ECMWF Report on ERS-2 RA for January 2010




Figure 10. Comparison of buoy wind speed observations with ERS2 Altimeter wind speed data for January 2010 (global)

ECMWF Report on ERS-2 RA for January 2010




Figure 11. Comparison of buoy wind speed observations with ERS2 Altimeter wind speed data for January 2010 (n.hem.)

■ ECMWF Report on ERS-2 RA for January 2010 ■




Figure 12. Comparison of buoy wind speed observations with ERS2 Altimeter wind speed data for January 2010 (hawaii)

ECMWF Report on ERS-2 RA for January 2010




Figure 13: ERS-2 Altimeter wind speeds: Timeseries of bias (ERS-2 - model) and scatter index (SI)

▪ ECMWF Report on ERS-2 RA for January 2010 ▪




Figure 14: Distribution of the ERS-2 Altimeter wave heights after QC for January 2010

▪ ECMWF Report on ERS-2 RA for January 2010 ▪




Figure 15: Distribution of ERS-2 Altimeter wave heights after along track averaging for January 2010

■ ECMWF Report on ERS-2 RA for January 2010 ■




Figure 16: Global distribution of ECMWF wave heights for January 2010

■ ECMWF Report on ERS-2 RA for January 2010 ■




Figure 17. Comparison of ECMWF wave height results with ERS2 Altimeter wave height data for January 2010 (global)

■ ECMWF Report on ERS-2 RA for January 2010 ■




Figure 18. Comparison of ECMWF wave height results with ERS2 Altimeter wave height data for January 2010 (n.hem.)

■ ECMWF Report on ERS-2 RA for January 2010 ■




Figure 19. Comparison of ECMWF wave height results with ERS2 Altimeter wave height data for January 2010 (tropics)

■ ECMWF Report on ERS-2 RA for January 2010 ■




Figure 20. Comparison of ECMWF wave height results with ERS2 Altimeter wave height data for January 2010 (s.hem.)

■ ECMWF Report on ERS-2 RA for January 2010 ■




Figure 21. Comparison of buoy wave height observations with ERS2 Altimeter wave height data for January 2010 (global)

■ ECMWF Report on ERS-2 RA for January 2010 ■




Figure 22. Comparison of buoy wave height observations with ERS2 Altimeter wave height data for January 2010 (n.hem.)

■ ECMWF Report on ERS-2 RA for January 2010 ■




Figure 23. Comparison of buoy wave height observations with ERS2 Altimeter wave height data for January 2010 (hawaii)

STATISTICS

ENTRIES	0
MEAN BUOY	0.0000
MEAN ERS-2	0.0000
BIAS (ERS-2 - BUOY)	0.0000
STANDARD DEVIATION	0.0000
SCATTER INDEX	0.0000
CORRELATION	0.0000
SYMMETRIC SLOPE	0.0000
REGR. COEFFICIENT	0.0000
REGR. CONSTANT	0.0000

ECMWF Report on ERS-2 RA for January 2010




Figure 24: ERS-2 Altimeter wave heights: Timeseries of bias (ERS-2 - model) and scatter index (SI)

ECMWF Report on ERS-2 RA for January 2010




Figure 25: ERS-2 Altimeter wave heights: Timeseries of bias (ERS-2 - model) and scatter index (SI)

ECMWF Report on ERS-2 RA for January 2010




Figure 26: ERS-2 Altimeter wind speeds: Timeseries of bias (ERS-2 - model) and scatter index (SI)

ECMWF Report on ERS-2 RA for January 2010




Figure 27: Significant wave height: Monthly mean difference of ERS-2 altimeter data minus wave model results for January 2010.