

■ ECMWF Report on ERS-2 RA for July 2006 ■

Title: Report on ERS-2 Radar Altimeter wave height and wind speed data.

By: *Saleh Abdalla*

Date: *4 August 2006*

Overview:

Based on the data received at ECMWF during the whole month, on average, 5226 observations arrived at ECMWF every 6 hours of which 68.23% passed the quality control. As can be seen in Figure 1, there was no data during the 6-hour time windows centred at 18:00 UTC on the 30th. and 00:00 and 06:00 UTC on the 31st. of the month. Furthermore, there was significant reduction of data at several time windows. The current coverage of the data covers: the North Atlantic, the western coast of North America, the eastern coast of China and around Korea and Japan, and the Southern Ocean as can be seen in Figure 27. The quality of the received data is as good as usual.

Backscatter:

ERS-2 $\langle\sigma_0\rangle = 11.73$ dB (with three peaks at 10.6 dB, 10.9 dB and 11.6 dB).

Wind Speed Comparison with ECMWF wind speeds (bias):

ERS-2 global: - 0.641 m/s

ERS-2 northern hemisphere: - 0.881 m/s

ERS-2 tropics: + 0.026 m/s

ERS-2 southern hemisphere: + 0.372 m/s

■ ECMWF Report on ERS-2 RA for July 2006 ■

Wind Speed Comparison with buoy wind speeds (bias):

ERS-2 global: - 1.108 m/s

ERS-2 northern hemisphere: - 1.109 m/s

ERS-2 tropics (Hawaii): *missing*

Wave Height Comparison with ECMWF wave heights (bias):

ERS-2 global: + 0.025 m (lowest waves measured: 0.6m)

ERS-2 northern hemisphere: +0.064 m

ERS-2 tropics: -0.008 m

ERS-2 southern hemisphere: - 0.240 m

Wave Height Comparison with buoy wave heights (bias):

ERS-2 global: + 0.089 m

ERS-2 northern hemisphere: + 0.086 m

ERS-2 tropics (Hawaii): *missing*

Remarks:

- The quality of Altimeter data, both wind speed and significant wave height, is as good as it used to be.
- Note that the large bias in both significant wave height and wind speed in the Southern Hemispheric extra tropics is due to the limited amount and coverage of the data.

■ ECMWF Report on ERS-2 RA for July 2006 ■

- According to ESA (EOHelp message of 4 July 2003), the recording capabilities of ERS-2 are declared permanently unavailable following the failure of the ERS-2 tape recorder on 22 June 2003. The ERS-2 tape recorders were used to record the ERS-2 Low Rate mission globally for a period of 8 years of continuous acquisition. The global coverage is, therefore, discontinued. The ERS-2 Low Rate mission will be continued within the visibility of ESA ground stations over Europe, North Atlantic, the Arctic and western North America. Coverage extended to include eastern coast of China and around Korea and Japan since 25 June 2005 and the Southern Ocean since 4 July 2005. Current coverage can be seen in Figure 27.
- Assimilation of ERS-2 RA wave heights into ECMWF wave model was stopped on 21st. of October 2003 and replaced by ENVISAT RA-2 Ku-band wave heights. Both instruments can not be assimilated at the same time as both satellites follow the same track with relatively short time separation (both can be assimilated safely if they were not so close).
- ENVISAT Ku-Band and Jason altimeter significant wave height data and ENVISAT ASAR Wave Mode Level 1b data are assimilated in the ECMWF wave model.
- There was no related ECMWF model changes this month (current operational cycle is CY30R1 since 1 February 2006).

Comparison Method:

The Altimeter wave height and wind speed data, as received by ECMWF from ESA through GTS, are the so-called fast delivery products. At ECMWF these data are subject to a quality control method, the details of which are described by Janssen et al. (1989) and Bauer et al. (1992). Consequently, superobservations are formed by averaging 30 consecutive data in order to match the spatial scales of the operational WAM model. Therefore, the collocation statistics are based on the comparison between these superobservations and operational wavemodel products.

■ ECMWF Report on ERS-2 RA for July 2006 ■

In addition, since also wave observations from buoys are received through the GTS, the Altimeter products are also compared against buoy observations. Again, in order to have matching scales, the buoy observations are averaged over a six hour time window. Apart from this, also a height correction is applied to the wind speed observations, since not all buoys observe the winds at the standard height of 10 m. A default observation height of 5 m is assumed, and when available the actual observation height is used. In order to interpolate from the observation height to the standard height a logarithmic wind profile with a roughness length as given by the Charnock relation is assumed, where the Charnock parameter is given the constant value of 0.018.

Figure captions:

- Figure 1: Time series of data reception for ERS-2 Altimeter data for July 2006.
- Figure 2: Distribution of the ERS-2 Altimeter Backscatter after QC for July 2006.
- Figure 3: Distribution of the ERS-2 Altimeter wind speeds after QC for July 2006.
- Figure 4: Distribution of the ERS-2 Altimeter wind speeds after along track averaging for July 2006.
- Figure 5: Global distribution of ECMWF ocean surface wind speeds for July 2006.
- Figure 6: Comparison of ECMWF wind speed results with ERS-2 Altimeter wind speed data for July 2006 (global).
- Figure 7: Comparison of ECMWF wind speed results with ERS-2 Altimeter wind speed data for July 2006 (northern hemisphere)
- Figure 8: Comparison of ECMWF wind speed results with ERS-2 Altimeter wind speed data for July 2006 (tropics)
- Figure 9: Comparison of ECMWF wind speed results with ERS-2 Altimeter wind speed data for July 2006 (southern hemisphere)
- Figure 10: Comparison of buoy wind speed observations with ERS-2 Altimeter wind speed data for July 2006 (global).
- Figure 11: Comparison of buoy wind speed observations with ERS-2 Altimeter wind speed data for July 2006 (northern hemisphere).
- Figure 12: Comparison of buoy wind speed observations with ERS-2 Altimeter wind speed data for July 2006 (hawaii).
- Figure 13: ERS-2 Altimeter wind speeds: Timeseries of bias (ERS-2 - model) and scatter index (SI).
- Figure 14: Distribution of the ERS-2 Altimeter wave heights after QC for July 2006.

■ ECMWF Report on ERS-2 RA for July 2006 ■

- Figure 15: Distribution of the ERS-2 Altimeter wave heights after along track averaging for July 2006.
- Figure 16: Global distribution of ECMWF wave heights for July 2006.
- Figure 17: Comparison of ECMWF wave height results with ERS-2 Altimeter wave height data for July 2006 (global).
- Figure 18: Comparison of ECMWF wave height results with ERS-2 Altimeter wave height data for July 2006 (northern hemisphere)
- Figure 19: Comparison of ECMWF wave height results with ERS-2 Altimeter wave height data for July 2006 (tropics)
- Figure 20: Comparison of ECMWF wave height results with ERS-2 Altimeter wave height data for July 2006 (southern hemisphere)
- Figure 21: Comparison of buoy wave height observations with ERS-2 Altimeter wave height data for July 2006 (global).
- Figure 22: Comparison of buoy wave height observations with ERS-2 Altimeter wave height data for July 2006 (northern hemisphere).
- Figure 23: Comparison of buoy wave height observations with ERS-2 Altimeter wave height data for July 2006 (hawaii).
- Figure 24: ERS-2 Altimeter wave heights: Timeseries of bias (ERS-2 - model) and scatter index (SI) for July 2006.
- Figure 25: ERS-2 Altimeter wave heights: Timeseries of bias (ERS-2 - model) and scatter index (SI) from December 1996 to July 2006.
- Figure 26: ERS-2 Altimeter wind speeds: Timeseries of bias (ERS-2 - model) and scatter index (SI) from December 1996 to July 2006.
- Figure 27: Significant wave height: Monthly mean difference of ERS-2 altimeter data minus wave model results for July 2006.

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 1: Time series of data reception for ERS-2 Altimeter data for July 2006

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 2: Distribution of the ERS-2 Altimeter backscatter after QC for July 2006

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 3: Distribution of the ERS-2 Altimeter wind speeds after QC for July 2006

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 4: Distribution of ERS-2 Altimeter wind speeds after along track averaging for July 2006

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 5: Global distribution of ECMWF ocean surface wind speeds for July 2006

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 6. Comparison of ECMWF wind speed results with ERS2 Altimeter wind speed data for July 2006 (global)

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 7. Comparison of ECMWF wind speed results with ERS2 Altimeter wind speed data for July 2006 (n.hem.)

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 8. Comparison of ECMWF wind speed results with ERS2 Altimeter wind speed data for July 2006 (tropics)

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 9. Comparison of ECMWF wind speed results with ERS2 Altimeter wind speed data for July 2006 (s.hem.)

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 10. Comparison of buoy wind speed observations with ERS2 Altimeter wind speed data for July 2006 (global)

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 11. Comparison of buoy wind speed observations with ERS2 Altimeter wind speed data for July 2006 (n.hem.)

▪ ECMWF Report on ERS-2 RA for July 2006 ▪




Figure 12. Comparison of buoy wind speed observations with ERS2 Altimeter wind speed data for July 2006 (hawaii)

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 13: ERS-2 Altimeter wind speeds: Timeseries of bias (ERS-2 - model) and scatter index (SI)

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 14: Distribution of the ERS-2 Altimeter wave heights after QC for July 2006

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 15: Distribution of ERS-2 Altimeter wave heights after along track averaging for July 2006

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 16: Global distribution of ECMWF wave heights for July 2006

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 17. Comparison of ECMWF wave height results with ERS2 Altimeter wave height data for July 2006 (global)

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 18. Comparison of ECMWF wave height results with ERS2 Altimeter wave height data for July 2006 (n.hem.)

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 19. Comparison of ECMWF wave height results with ERS2 Altimeter wave height data for July 2006 (tropics)

▪ ECMWF Report on ERS-2 RA for July 2006 ▪




Figure 20. Comparison of ECMWF wave height results with ERS2 Altimeter wave height data for July 2006 (s.hem.)

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 21. Comparison of buoy wave height observations with ERS2 Altimeter wave height data for July 2006 (global)

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 22. Comparison of buoy wave height observations with ERS2 Altimeter wave height data for July 2006 (n.hem.)

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 23. Comparison of buoy wave height observations with ERS2 Altimeter wave height data for July 2006 (hawaii)

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 24: ERS-2 Altimeter wave heights: Timeseries of bias (ERS-2 - model) and scatter index (SI)

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 25: ERS-2 Altimeter wave heights: Timeseries of bias (ERS-2 - model) and scatter index (SI)

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 26: ERS-2 Altimeter wind speeds: Timeseries of bias (ERS-2 - model) and scatter index (SI)

■ ECMWF Report on ERS-2 RA for July 2006 ■




Figure 27: Significant wave height: Monthly mean difference of ERS-2 altimeter data minus wave model results for July 2006.