

Monitoring statistics of the ERS-2 scatterometer for ESA cycle 106

(Project Ref. 18212/04/I-OL)

Hans Hersbach

European Centre for Medium-Range Weather Forecasts,

Shinfield Park, Reading, RG2 9AX, England

Tel: (+44 118) 9499476, e-mail: dal@ecmwf.int

July 15, 2005

1 Introduction

The quality of the UWI product was monitored at ECMWF for cycle 106. Results were compared to those obtained from the previous cycle, as well for data received during the nominal period in 2000 (up to cycle 59). No corrections for duplicate observations were applied.

During cycle 106 data was received between 21:02 UTC 6 June 2005 and 19:46 UTC 11 July 2005. No data was received for the 6-hourly batch for 00 UTC and 06 UTC 21 June 2005, and 18 UTC 29 June 2005.

Data is being recorded whenever within the visibility range of a ground station. In addition to the existing list of stations, data was received from McMurdo (Antarctica) and Beijing, from 01:24 UTC 25 June 2005 and 13:34 UTC 5 July 2005, respectively, onwards. Data volume from these newly included stations is, for the time being, low. For cycle 106 data coverage was found to extend over the North-Atlantic, part of the Mediterranean, the Caribbean, the Gulf of Mexico, a small part of the Pacific west from the US Canada and Central America, the Chinese and Japanese Sea, and the Southern Ocean south of Australia and New Zealand (see Figure 2).

The asymmetry between the fore and aft incidence angles showed many peaks, mostly behaving within reasonable bounds. The k_p -yaw ESA flag set accordingly. Between 6 and 8 July 2005, an enhanced volatility was observed. It did not coincide with high solar activity (source: www.spaceweather.com).

Compared to cycle 105, the UWI wind speed relative to ECMWF first-guess (FG) fields showed a record-low relative standard deviation (from 1.35 m/s to 1.29

m/s). However, the relative bias grew to exceptionally large negative values (from -0.80 m/s to -1.02 m/s), and similarly for CMOD5 winds (from -0.32 m/s to -0.59 m/s). Standard deviation of the UWI winds is lower than that for 2000; the bias is more negative (Figure 1). The situation is similar to that encountered during Summer 2004.

The performance of the UWI wind direction was stable during cycle 106.

Ocean calibration shows an increased inter-node and inter-beam dependency of bias levels. The average bias level became significantly more negative (-0.91 dB, was -0.58 dB), especially at high incidence angles (figure 4).

The ECMWF assimilation/forecast system was changed on 28 June 2005. The assimilation of SSM/I and AIRS radiances was adapted, Meteosat-8 (MSG) winds were included, surface-pressure bias corrections were refined, the estimation for the background error was changed, and the SMHI Baltic sea-ice analysis was included. Model refinements included reduction of spin-up effects and changes in convection. The anticipated effect on surface winds is limited.

The cycle-averaged evolution of performance relative to ECMWF first-guess (FG) winds is displayed in Figure 1. Figure 2 shows global maps of the over cycle 106 averaged UWI data coverage and wind climate, Figure 3 for performance relative to FG winds.

2 ERS-2 statistics from 7 June to 11 July 2005

2.1 Sigma0 bias levels

The average sigma0 bias levels (compared to simulated sigma0's based on ECMWF model FG winds) stratified with respect to antenna beam, ascending or descending track and as function of incidence angle (i.e. across-node number) is displayed in Figure 4.

Inter-node and inter-beam dependencies have grown considerably. Bias levels of the fore and aft beam are the most negative ones, especially at high incidence angles. Average bias level has grown from -0.58 dB to -0.91 dB, being 0.2 dB more negative than for nominal data in 2000 (see Figure 1 of the reports for cycle 48 to 59). The situation is very similar to that of one year ago (see report for cycle 96), where large negative biases in both wind and backscatter were observed.

The data volume of descending tracks was about 10% higher than for ascending tracks.

2.2 Incidence angles

For ESACA, across-node binning is, like the old processor, retained on a 25km mesh. From simple geometrical arguments it follows that variations in yaw attitude will lead to asymmetries between the incidence angles of the fore and aft beam. Indeed, this has been observed. Figure 5 gives a time evolution of this asymmetry, showing rapid variations, which are typical for yaw attitude errors. Also in this figure, the

occasions for which the combined k_p -yaw quality flag was set are indicated by red stars. The relation with incidence-angle asymmetries is obvious.

During cycle 106 most peaks were within bounds. Between 6 and 8 July 2005 a period with high volatility was observed, during which solar activity was normal (source www.spaceweather.com).

2.3 Distance to cone history

The distance to the cone history is shown in Figure 6. Curves are based on data that passed all QC, including the test on the k_p -yaw flag, and subject to the land and sea-ice check at ECMWF (see cyclic report 88 for details).

Like for cycle 105, time series are (due to lack of statistics) very noisy, especially for the near-range nodes. Most spikes were found to be the result of low data volumes.

Compared to cycle 105, the average level was slightly higher (from 1.22 to 1.18), and is now about 12% higher than for nominal data (see top panel Figure 1). Since the end of June 2005, levels have seemed to increase.

2.4 UWI minus First-Guess wind history

In Figure 7, the UWI minus ECMWF first-guess wind-speed history is plotted.

The history plot shows several peaks, most of which are related to low data volumes. Similar results apply for the history of de-aliased CMOD4 winds versus FG (Figure 9).

Figure 11 displays the locations for which UWI winds were more than 8 m/s weaker (top panel) and more than 8 m/s stronger (lower panel) than FG winds. Like for cycle 105, the number of such collocations is low. Two cases for stronger UWI winds are presented in Figure 12. The top panel shows the capture of Hurricane Dennis on 7 July 2005, just before it made landfall over Cuba, leaving a trail of destruction. The (CMOD5) winds show, like the ECMWF first-guess, a detailed structure, however more intense than the latter. Note the erroneous looking patch in the bottom of the panel.

The lower panel of Figure 12 shows a case from the newly included station at McMurdo. Although the strong UWI and ECMWF winds for this winter case in the Southern Hemisphere storm track may locally differ significantly, the flow of both fields looks, modulo de-aliasing errors, sensible. In Figure 3 the quality of UWI winds may appear lower than average for this new station. However, the newly covered region is characterised by strong winds with high variability, which, given the low data volume make an objective comparison difficult.

Average bias levels and standard deviations of UWI winds relative to FG winds are displayed in Table 1. From this it is seen that the bias of both the UWI and CMOD4 product have become more negative. The average bias level is lower to that for nominal data in 2000 (UWI: -1.02 m/s now, was -0.79 m/s for cycle 59). The evolution of the bias from cycles 92 to 106 is displayed in the top panel of Figure 17. The red curve is the 15-day moving average for the at ECMWF inverted

	cycle 105		cycle 106	
	UWI	CMOD4	UWI	CMOD4
speed STDV	1.35	1.34	1.29	1.29
node 1-2	1.39	1.37	1.35	1.35
node 3-4	1.33	1.33	1.30	1.30
node 5-7	1.30	1.30	1.25	1.26
node 8-10	1.31	1.31	1.25	1.26
node 11-14	1.31	1.31	1.24	1.25
node 15-19	1.35	1.35	1.28	1.28
speed BIAS	-0.80	-0.80	-1.02	-1.02
node 1-2	-1.27	-1.25	-1.40	-1.39
node 3-4	-1.04	-1.00	-1.21	-1.18
node 5-7	-0.81	-0.79	-1.04	-1.02
node 8-10	-0.64	-0.64	-0.91	-0.91
node 11-14	-0.64	-0.66	-0.87	-0.89
node 15-19	-0.66	-0.69	-0.89	-0.92
direction STDV	26.2	18.8	27.2	18.4
direction BIAS	-2.7	-2.7	-3.3	-3.7

Table 1: Biases and standard deviation of ERS-2 versus ECMWF FG winds in m/s for speed and degrees for direction.

ERS-2 winds; i.e., CMOD5 since 9 March 2004. Blue vertical dashed lines indicate ECMWF model changes. This plot shows that the rapidly evolving negative bias for ERS-2 starting in March 2005, temporarily stabilizing in May 2005, and a fast drop again during the last two months. The situation of such large negative biases is similar to that of the summer of 2004. For the global average of QuikSCAT winds (lower panel of Figure 17), there is an indication of a similar, though much smaller trend.

The standard deviation of UWI wind speed compared to cycle 105 has been reduced significantly (1.29 m/s, was 1.35 m/s), the main reason being a milder wind climate.

For cycle 106 the (UWI - FG) direction standard deviations were mostly ranging between 15 and 40 degrees (Figure 8). Sharp peaks are the result of low data volumes. The average performance for UWI wind direction was slightly lower (STDV 27.2 degrees, was 26.2 degrees, bias -3.3 degrees, was -2.7 degrees), that of de-aliased CMOD4 winds hardly changed (18.4 degrees, was 18.8 degrees).

2.5 Scatterplots

Scatterplots of FG winds versus ERS-2 winds are displayed in Figures 13 to 16. Values of standard deviations and biases are slightly different from those displayed in Table 1. Reason for this is that, for plotting purposes, the in 0.5 m/s resolution ERS-2 winds have been slightly perturbed (increases scatter with 0.02 m/s), and that zero wind-speed ERS-2 winds have been excluded (decreases scatter with about

0.05 m/s).

The scatterplot of UWI wind speed versus FG (Figure 13) is very similar to that for (at ECMWF inverted) de-aliased CMOD4 winds (Figure 15). It confirms that the ESACA inversion scheme is working properly.

Winds derived on the basis of CMOD5 are displayed in Figure 16. The relative standard deviation is lower than for CMOD4 winds (1.29 m/s versus 1.31 m/s). Compared to ECMWF FG, CMOD5 winds are -0.59 m/s slower; this average arising from mostly moderate winds. The agreement for the more extreme winds, fortunately, looks fine.

Figure Captions

Figure 1: Evolution of the performance of the ERS-2 scatterometer averaged over 5-weekly cycles from 12 December 2001 (cycle 69) to 11 July 2005 (end cycle 106) for the UWI product (solid, star) and de-aliased winds based on CMOD4 (dashed, diamond). Results are based on data that passed the UWI QC flags. For cycle 85 two values are plotted; the first value for the global set, the second one for the regional set. Dotted lines represent values for cycle 59 (5 December 2000 to 17 January 2001), i.e. the last stable cycle of the nominal period. From top to bottom panel are shown the normalized distance to the cone (CMOD4 only) the standard deviation of the wind speed compared to FG winds, the corresponding bias (for UWI winds the extreme inter-node averages are shown as well), and the standard deviation of wind direction compared to FG.

Figure 2: Average number of observations per 12H and per 125km grid box (top panel) and wind-climate (lower panel) for UWI winds that passed the UWI flags QC and a check on the collocated ECMWF land and sea-ice mask.

Figure 3: The same as Figure 2, but now for the relative bias (top panel) and standard deviation (lower panel) with ECMWF first-guess winds.

Figure 4: Ratio of $< \sigma_0^{0.625} > / < \text{CMOD4}(\text{FirstGuess})^{0.625} >$ converted in dB for the fore beam (solid line), mid beam (dashed line) and aft beam (dotted line), as a function of incidence angle for descending and ascending tracks. The thin lines indicate the error bars on the estimated mean. First-guess winds are based on the in time closest (+3h, +6h, +9h, or +12h) T511 forecast field, and are bilinearly interpolated in space.

Figure 5: Time series of the difference in incidence angle between the fore and aft beam. Red stars indicate the occurrences for which the combined k_p -yaw flag was set.

Figure 6: Mean normalized distance to the cone computed every 6 hours for nodes 1-2, 3-4, 5-7, 8-10, 11-14 and 15-19 (solid curve close to 1 when no instrumental problems are present). The dotted curve shows the number of incoming triplets in logarithmic scale (1 corresponds to 60,000 triplets) and the dashed one indicates the fraction of complete (based on the land and sea-ice mask at ECMWF) sea-located triplets rejected by ESA flags, or by the wind inversion algorithm (0: all data kept, 1: no data kept).

Figure 7: Mean (solid line) and standard deviation (dashed line) of the wind speed difference UWI - first guess for the data retained by the quality control.

Figure 8: Same as Fig. 7, but for the wind direction difference. Statistics are computed only for wind speeds higher than 4 m/s.

Figures 9 and 10: Same as Fig. 7 and 8 respectively, but for the de-aliased CMOD4 data.

Figure 11: Locations of data during cycle 106 for which UWI winds are more than 8 m/s weaker (top panel) respectively stronger (lower panel) than FG, and on which QC on UWI flags and the ECMWF land/sea-ice mask was applied.

Figure 12: Comparison between UWI (red) and ECMWF FG (blue) winds for hurricane Dennis on 7 July 2005 (top panel) and for a case in the Southern-Hemisphere storm track, south of New Zealand on 10 July 2005 (lower panel).

Figure 13: Two-dimensional histogram of first guess and UWI wind speeds, for the data kept by the UWI flags, and QC based on the ECMWF ice and land and sea-ice mask. Circles denote the mean values in the y-direction, and squares those in the x-direction.

Figure 14: Same as Fig. 13, but for wind direction. Only wind speeds higher than 4m/s are taken into account.

Figure 15: Same as Fig. 13, but for de-aliased CMOD4 winds.

Figure 16: Same as Fig. 13, but for de-aliased CMOD5 winds.

Figure 17: Bias relative to FG winds of the wind speed of ERS-2 winds (based on bias-corrected CMOD4 before 9 March 2004, and on CMOD5 afterwards) for nodes 1-19 (top panel) respectively of 50-km QuikSCAT (based on the QSCAT-1 model function) for nodes 5-34 (i.e., inner-beam zone; middle and lower panels) versus ECMWF first guess for the period of cycle 92 to 106. Curves represent centred 15-day running means. Vertical dashed blue lines mark ECMWF model changes.





Figure 1

NOBS (ERS-2 UWI), per 12H, per 125km box
average from 2005060700 to 2005071118 GLOB:3.04



AVERAGE (ERS-2 UWI), in m/s.
average from 2005060700 to 2005071118 GLOB:5.35





Figure 2

BIAS (ERS-2 UWI vs FIRST-GUESS), in m/s.
average from 2005060700 to 2005071118 GLOB:-1.07



STDV (ERS-2 UWI vs FIRST-GUESS), in m/s.
average from 2005060700 to 2005071118 GLOB:1.08




Figure 3





Figure 4




Figure 5

Monitoring of Sigma0 triplets versus CMOD4 for ERS-2

from 2005060700 to 2005071118

(solid) mean normalised distance to the cone over 6 h

(dashed) fraction of complete sea-point observations rejected by ESA flag or CMOD4 inversion

(dotted) total number of data in log. scale (1 for 60000)










Figure 6

Monitoring of UWI winds versus First Guess for ERS-2

from 2005060700 to 2005071118

(solid) wind speed bias UWI - First Guess over 6h (deg.)

(dashed) wind speed standard deviation UWI - First Guess over 6h (deg.)



2005

2005

Figure 7

Monitoring of UWI winds versus First Guess for ERS-2

from 2005060700 to 2005071118

(solid) wind direction bias UWI - First Guess over 6h (deg.)

(dashed) wind direction standard deviation UWI - First Guess over 6h (deg.)















Figure 8

Monitoring of de-aliased CMOD4 winds versus First Guess for ERS-2

from 2005060700 to 2005071118

(solid) wind speed bias CMOD4 - First Guess over 6h (deg.)

(dashed) wind speed standard deviation CMOD4 - First Guess over 6h (deg.)



2005

2005







Figure 9

Monitoring of de-aliased CMOD4 winds versus First Guess for ERS-2

from 2005060700 to 2005071118

(solid) wind direction bias CMOD4 - First Guess over 6h (deg.)

(dashed) wind direction standard deviation CMOD4 - First Guess over 6h (deg.)




2005

2005

Figure 10

UWI winds more than 8 m/s weaker than FGAT

CYCLE 106, 2005060700 to 2005071118, QC on ESA flags



UWI winds more than 8 m/s stronger than FGAT

CYCLE 106, 2005060700 to 2005071118, QC on ESA flags





Figure 11

CMOD5 winds (red) versus FGAT winds (blue)
20050707 15:16 UTC



CMOD4 winds (red) versus FGAT winds (blue)
20050710 22:26 UTC




Figure 12




Figure 13

ECMWF 3-hourly First-Guess winds versus UWI winds
from 2005060700 to 2005071118
= 625032 (|f| gt 4.00 m/s), db contour levels, 5 db step, 1st level at 3.0 db
m(y-x)=-3.58 sd(y-x)= 27.19 sdx=102.10 sdy=101.77 pcxy= 0.982




Figure 14




Figure 15





Figure 16

ERS-2 (CMOD5)



QuikSCAT (QSCAT-1)




Figure 17