


Biotoxicological risks and hazards of a thawing Arctic



Dr. Kimberley R. Miner
NASA Jet Propulsion Laboratory
California Institute of Technology



Age of Permafrost

- Permafrost remaining from the last ice age (c. 115,000 – c. 11,700 years ago) and prior glaciations
- Contains organic carbon -plant and animal detritus, leads to carbon release in the Permafrost Carbon Feedback
- Rapid thaw in the last 50 years is changing structural dynamics of the tundra, releasing GHG and permafrost components
- Thaw dynamics based upon underlying structure of the permafrost and temperature

Miner et al. 2021 *Nature Climate Change*, Solicited review



Yedoma and Thermokarst

- Yedoma can be 100,000 of years old and hundreds of meters deep- exposed by thermokarst development
- May contain viable microbes from the Cenozoic to modern times
- Known microbes include methanogenic microorganisms, responsible for releasing CH₄ and CO₂
- 2% carbon by mass, but 80-90% ice
- Potential for a 50% increase in released carbon when exposed to the atmosphere




Permafrost microbes



Permafrost microbes

- Cold-adapted generalist microbes with lipid membranes still viable (extremophiles)
 - Cold methane seeps: ancient bacteria phylotypes related to *Loktanella*, *Gillisia*, *Halomonas*, and *Marinobacter* spp.
 - Brine lenses (100k-120kyr): *Clostridium* and *Psychrobacter*
 - Permafrost substrate (30kyr): intact virus species including Mimiviruses, Pandoraviruses, *Pithovirus sibericum*, *Mollivirus sibericum*
- Unknown bacteria, viruses and microbes
 - Up to 1 million years old
 - ‘Methuselah’ microbes



Permafrost microbes

- The Plague, Black Death, Small Pox and unknown viruses including new "Alaskapox"
- Antibiotic resistance in some species
 - bacteria resistant to chloramphenicol, streptomycin, kanamycin, gentamicin, tetracycline, spectinomycin, and neomycin have been recovered (15-290kyr)
- Other potential 'eradicated' diseases



Potential Impacts

- Anthrax release in Siberia suspected in death of 200 reindeer and a child in 2016
- Graves of Smallpox victims excavated in 2016, 2019
- Microbes from 700-yr old Caribou cloned into modern plant in 2014
- Exploration of microbiome in unprotected labs ongoing
- Exposure can expand with tourism travel to the Arctic



Anthropogenic contamination



Persistent Organic Pollutants (POPs)

- Organic chemicals including DDT, HCH, PCBs deposited atmospherically
 - 2019, 2020 studies show human risk from fish consumption
 - High concentrations- DDT (10 ng/L) , PCB (~4.5 ng/L), and HCH (~20 ng/L) in glacier ice and permafrost
 - Plastics, fragrances, and plasticizers (PAHs, PFAS, brominated flame retardants)



Heavy Metals

- Heavy metals Arsenic (As), Cd, Nickel (Ni), and Mercury (Hg) from mine tailings and atmospheric deposition
- One open pit mine: Cd (1 – 4 mg/kg), Ni (1000 – 1500 mg/kg), and Hg (40-120 mg/kg)
- Estimated 384- 1,656 ± 962 Gg Hg in the top three meters of soil
- Bioaccumulation in fish may see an increase of up to 222% by 2300
- *Glaucous Gull* species show high levels of Hg (4.9 ug/g), PCB (3326 ng/g), and DDT (2367 ng/g)



Nuclear waste

- 1955-1990, the Soviet Union conducted 130 nuclear weapons tests in the Novaya Zemlya Archipelago
 - 224 separate explosive devices, releasing ~265 megatons of nuclear energy
 - Kara Sea sediments contain up to 11,000 Bq/kg of Plutonium, 3-4x larger than the background
 - Sunken ships excluded from cleanup account for ~8860 TBq of radiation
- Camp Century: radioactivity of $\sim 1.2 \times 10^9$ Bq including physical waste, diesel fuel & PCBs, and sewage
- 1968 Thule bomber crash $> 4.6 \times 10^{12}$ Bq of Uranium and Plutonium on Greenland ice sheet





Contaminant release

- Melt and thaw of ice and permafrost
- Release during industrial use (storage and seepage)
- Nuclear materials mined, utilized or stored in the Arctic
- Release from accidents, global transport or shipwrecks
- Oil drilling, transport and infrastructure leaks



Could Arctic hazards transport globally?

- Potential introduction of old bacteria, viruses, and microbial life into land and ocean
- Atmospheric remobilization of chemicals and toxins stored in the Arctic
- The 'greening' of the Arctic and changing hydrology
- A possible route for microbial contamination in freshwater



Next Steps


- NASA-ESA Arctic Methane Challenge
- Core sampling in permafrost tunnels, flow modeling in Greenland
- Combined satellite and in-situ monitoring
- Work with Mars Rover tools- linking planetary exploration and extremophiles




Next steps: microbes in permafrost and in space

- Goal: To find, characterize and index the permafrost microbiome, developing a greater understanding of niches supporting life
- Sampling permafrost from 10-130,000 years old in Alaska
- Proof of concept in Greenland & on the Mars Perseverance a similar instrument (SHERLOC) with same laser system
- **If we can identify ancient, extremophile organisms on Earth, successful recovery from similar extraplanetary environments may be possible**

Visible Image



Fluorescence



SHERLOC
Jet Propulsion Laboratory
California Institute of Technology

Questions?

Kimberley.n.miner@jpl.nasa.gov

drkimberleyrain.com



Figure: Victor Leshyk

1. Vincent, W. F., Lemay, M. & Allard, M. Arctic permafrost landscapes in transition: towards an integrated Earth system approach. *Arct. Sci.* **3**, 39–64 (2017).
2. Hjort, J. *et al.* Degrading permafrost puts Arctic infrastructure at risk by mid-century. *Nat. Commun.* **9**, (2018).
3. Karjalainen, O. *et al.* Data Descriptor: Circumpolar permafrost maps and geohazard indices for near-future infrastructure risk assessments Background & Summary. *Nat. Publ. Gr.* (2019) DOI:10.1038/sdata.2019.37.
4. Waits, A., Emelyanova, A., Oksanen, A., Abass, K. & Rautio, A. Human infectious diseases and the changing climate in the Arctic. *Environ. Int.* **121**, 703–713 (2018).
5. Miner, K. R. *et al.* A screening-level approach to quantifying risk from glacial release of organochlorine pollutants in the Alaskan Arctic. *J. Expo. Sci. Environ. Epidemiol.* (2018) doi:doi.org/10.1038/s41370-018-0100-7.
6. Colgan, W. *et al.* The abandoned ice sheet base at Camp Century, Greenland, in a warming climate. *Geophys. Res. Lett.* **43**, 8091–8096 (2016).
7. Eriksson, M., Lindahl, P., Roos, P., Dahlggaard, H. & Holm, E. U, Pu, and Am nuclear signatures of the thule hydrogen bomb debris. *Environ. Sci. Technol.* **42**, 4717–4722 (2008).
8. Perryman, C. R. *et al.* Heavy metals in the Arctic: Distribution and enrichment of five metals in Alaskan soils. *PLoS One* **15**, 1–14 (2020).
9. Revich, B., Tokarevich, N. & Parkinson, A. J. Climate change and zoonotic infections in the Russian arctic. *Int. J. Circumpolar Health* **71**, (2012).
10. Kaab, A. Remote sensing of permafrost-related problems and hazards. *Permafrost. Periglac. Process.* **136**, 107–136 (2008).
11. Rivkina, E. *et al.* Biogeochemistry of methane and methanogenic archaea in permafrost. *FEMS Microbiol. Ecol.* **61**, 1–15 (2007).
12. Zhang, D. C., Brouckov, A., Griva, G., Schinner, F. & Margesin, R. Isolation and characterization of bacteria from ancient Siberian permafrost sediment. *Biology (Basel)*. **2**, 85–106 (2013).
13. Wurzbacher, C., Nilsson, R. H., Rautio, M. & Peura, S. Poorly known microbial taxa dominate the microbiome of permafrost thaw ponds. *ISME J.* **11**, 1938–1941 (2017).
14. Schütte, U. M. E. *et al.* Effect of permafrost thaw on plant and soil fungal community in a boreal forest: Does fungal community change mediate plant productivity response? *J. Ecol.* **107**, 1737–1752 (2019).
15. Diversity, B. Arctic Ocean Drilling: Risking Oil Spills, Human Life, and Wildlife. (2012).
16. Goodman, S. & Kertysova, K. The nuclearisation of the Russian Arctic : new reactors , new risks POLICY BRIEF. *Eur. Leadersh. Netw.* (2020).
17. Byrne, S. *et al.* Persistent organochlorine pesticide exposure related to a formerly used defense site on St. Lawrence Island, Alaska: data from sentinel fish and human sera. *Toxicol. Environ. Heal.* **78**, 37–54 (2015).
18. Mackay, D. *Multimedia Environmental Models*. (CRC Press, 2001, 2001).
19. Abbott, B. W. *et al.* Biomass offsets little or none of permafrost carbon release from soils, streams, and wildfire: An expert assessment. *Environ. Res. Lett.* **11**, (2016).
20. Ren, J. *et al.* Biomagnification of persistent organic pollutants along a high-altitude aquatic food chain in the Tibetan Plateau: Processes and mechanisms. *Environ. Pollut.* 1–8 (2016) doi:10.1016/j.envpol.2016.10.019.
21. Jia, G. J., Epstein, H. E. & Walker, D. A. Vegetation greening in the Canadian Arctic related to decadal